




PCT/GB 2003 / 0 0 0 0 6 3



INVESTOR IN PEOPLE

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office

Concept House

Cardiff Road

Newport

South Wales

NP10 8QQ

REC'D 18 FEB 2003

PO


PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *AmBrewer*

Dated 24 January 2003


BEST AVAILABLE COPY

Patents Form 1/77

Patents Act 1977
(16)



21 JAN 02 EGB9152-1 002093
P01/7700 0.00-0201160.9



Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference **PPD 50680/GB/P**

2. Patent application number
(The Patent Office will fill in this part)

0201160.9

3. Full name, address and postcode of the or of each applicant (underline all surnames)

**SYNGENTA LIMITED
FERNHURST
HASLEMERE
SURREY
GU27 3JE
UNITED KINGDOM**

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

8038028001

4. Title of the invention

BIOCIDAL COMPOUND AND ITS PREPARATION

5. Name of your agent (if you have one)

Malcolm John HOUGHTON

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

**SYNGENTA Limited
Intellectual Property Dept
Jealott's Hill International Research Centre
PO Box No 3538
Bracknell
Berkshire RG43 6YA
United Kingdom**

Patents ADP number (if you know it)

8148298002

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country	Priority application number (if you know it)	Date of filing (day / month / year)
---------	---	--

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application	Date of filing (day / month / year)
-------------------------------	--

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

YES

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description	25	✓
Claim(s)	2	✓
Abstract		✓
Drawing(s)		✓

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature *J A Bowdich*
Julie Ann BOWDICH

Date 18 January 2001

12. Name and daytime telephone number of person to contact in the United Kingdom

JULIE Ann BOWDICH 01344 414102

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.


Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

BIOCIDAL COMPOUND AND ITS PREPARATION

This invention relates to a new biocidal compound and to a process for its preparation. More particularly it relates to a biocidal compound which may be obtained by cultivation of *Streptomyces* organisms; to the use of said compound as a biocide; to compositions containing a biocidally effective amount of said compound; a method of combating pathogens using said compounds; a new strain of *Streptomyces malaysiensis* and mutants thereof.


In a first aspect the invention provides a new biocidal compound of formula (I)



This is a novel compound not previously isolated. It is characterised by the nmr and mass spectroscopy data provided in the examples herein. It may be prepared by growing under controlled conditions, a previously unknown strain of micro-organism from the species *Streptomyces malaysiensis*. The compound of the invention has in particular, insecticidal, herbicidal, anti-fungal, anti-viral and anti-cancer activity and is of special interest for use in agriculture, horticulture, animal and human health. The compound may also be of use as an intermediate in the preparation of further active compounds. The compound of formula (I) may be obtained by fermentation and recovered in substantially pure form as described herein.

The compound of formula (I) has six asymmetric centres and may exist in the form of one or more isomers. The invention includes all isomers, individually or in a mixture of one or more forms. The compound of the invention may exist in several tautomeric forms and the invention extends to all forms.

Of particular interest is the compound of formula (II)



The compound of formula (II) may exist in alternative tautomeric forms and the invention extends to said alternative forms.

The compound of formula (I) is an active antiviral agent showing for example
 5 antiviral activity against the Human Immunodeficiency Virus associated with the disease AIDS. The compound of formula (I) has also been shown to have anti-cancer properties, for example it shows an inhibitory effect on the cell proliferation of cancer cells. As used herein the term biocidal is used to cover activity against pathogens and includes antiviral, antifungal and anticancer activity. The use of the compound as an antifungal agent
 10 particularly against plant pathogenic fungi is particularly preferred according to the invention.

The compound of formula (I) is an active fungicide and may be used to control one or more of the following plant pathogens: *Pyricularia oryzae* (*Magnaporthe grisea*) on rice and wheat and other *Pyricularia* spp. on other hosts; *Puccinia triticina* (or *recondita*), *Puccinia striiformis* and other rusts on wheat, *Puccinia hordei*, *Puccinia striiformis* and other rusts on
 15 barley, and rusts on other hosts (for example turf, rye, coffee, pears, apples, peanuts, sugar beet, vegetables and ornamental plants); *Erysiphe cichoracearum* on cucurbits (for example melon); *Blumeria* (or *Erysiphe*) *graminis* (powdery mildew) on barley, wheat, rye and turf and other powdery mildews on various hosts, such as *Sphaerotheca macularis* on hops,
 20 *Sphaerotheca fusca* (*Sphaerotheca fuliginea*) on cucurbits (for example cucumber), *Leveillula taurica* on tomatoes, aubergine and green pepper, *Podosphaera leucotricha* on apples and *Uncinula necator* on vines; *Cochliobolus* spp., *Helminthosporium* spp., *Drechslera* spp. (*Pyrenophora* spp.), *Rhynchosporium* spp., *Mycosphaerella graminicola* (*Septoria tritici*) and *Phaeosphaeria nodorum* (*Stagonospora nodorum* or *Septoria nodorum*), *Pseudocercospora herpotrichoides* and *Gaeumannomyces graminis* on cereals
 25 (for example wheat, barley, rye), turf and other hosts; *Cercospora arachidicola* and *Cercosporidium personatum* on peanuts and other *Cercospora* spp. on other hosts, for example sugar beet, bananas, soya beans and rice; *Botrytis cinerea* (grey mould) on

tomatoes, strawberries, vegetables, vines and other hosts and other *Botrytis* spp. on other hosts; *Alternaria* spp. on vegetables (for example carrots), oil-seed rape, apples, tomatoes, potatoes, cereals (for example wheat) and other hosts; *Venturia* spp. (including *Venturia inaequalis* (scab)) on apples, pears, stone fruit, tree nuts and other hosts; *Cladosporium* spp. on a range of hosts including cereals (for example wheat) and tomatoes; *Monilinia* spp. on stone fruit, tree nuts and other hosts; *Didymella* spp. on tomatoes, turf, wheat, cucurbits and other hosts; *Phoma* spp. on oil-seed rape, turf, rice, potatoes, wheat and other hosts; *Aspergillus* spp. and *Aureobasidium* spp. on wheat, lumber and other hosts; *Ascochyta* spp. on peas, wheat, barley and other hosts; *Stemphylium* spp. (*Pleospora* spp.) on apples, pears, onions and other hosts; summer diseases (for example bitter rot (*Glomerella cingulata*), black rot or frog-eye leaf spot (*Botryosphaeria obtusa*), Brooks fruit spot (*Mycosphaerella pomi*), Cedar apple rust (*Gymnosporangium juniperi-virginianae*), sooty blotch (*Gloeodes pomigena*), flyspeck (*Schizothyrium pomi*) and white rot (*Botryosphaeria dothidea*)) on apples and pears; *Plasmopara viticola* on vines; other downy mildews, such as *Bremia lactucae* on lettuce, *Peronospora* spp. on soybeans, tobacco, onions and other hosts, *Pseudoperonospora humuli* on hops and *Pseudoperonospora cubensis* on cucurbits; *Pythium* spp. (including *Pythium ultimum*) on turf and other hosts; *Phytophthora infestans* on potatoes and tomatoes and other *Phytophthora* spp. on vegetables, strawberries, avocado, pepper, ornamentals, tobacco, cocoa and other hosts; *Thanatephorus cucumeris* on rice and turf and other *Rhizoctonia* spp. on various hosts such as wheat and barley, peanuts, vegetables, cotton and turf; *Sclerotinia* spp. on turf, peanuts, potatoes, oil-seed rape and other hosts; *Sclerotium* spp. on turf, peanuts and other hosts; *Gibberella fujikuroi* on rice; *Colletotrichum* spp. on a range of hosts including turf, coffee and vegetables; *Laetisaria fuciformis* on turf; *Mycosphaerella* spp. on bananas, peanuts, citrus, pecans, papaya and other hosts; *Diaporthe* spp. on citrus, soybean, melon, pears, lupin and other hosts; *Elsinoe* spp. on citrus, vines, olives, pecans, roses and other hosts; *Verticillium* spp. on a range of hosts including hops, potatoes and tomatoes; *Pyrenopeziza* spp. on oil-seed rape and other hosts; *Oncobasidium theobromae* on cocoa causing vascular streak dieback; *Fusarium* spp., *Typhula* spp., *Microdochium nivale*, *Ustilago* spp., *Urocystis* spp., *Tilletia* spp. and *Claviceps purpurea* on a variety of hosts but particularly wheat, barley, turf and maize; *Ramularia* spp. on sugar beet, barley and other hosts; post-harvest diseases particularly of fruit (for example *Penicillium digitatum*, *Penicillium italicum* and *Trichoderma viride* on oranges, *Colletotrichum musae* and *Gloeosporium musarum* on bananas and *Botrytis cinerea* on

grapes); other pathogens on vines, notably *Eutypa lata*, *Guignardia bidwellii*, *Phellinus igniarius*, *Phomopsis viticola*, *Pseudopeziza tracheiphila* and *Stereum hirsutum*; other pathogens on trees (for example *Lophodermium seeditiosum*) or lumber, notably *Cephaloscypha frugifera*, *Ceratocystis* spp., *Ophiostoma piceae*, *Penicillium* spp.,

5 *Trichoderma pseudokoningii*, *Trichoderma viride*, *Trichoderma harzianum*, *Aspergillus niger*, *Leptographium lindbergii* and *Aureobasidium pullulans*; and fungal vectors of viral diseases (for example *Polymyxa graminis* on cereals as the vector of barley yellow mosaic virus (BYMV) and *Polymyxa betae* on sugar beet as the vector of rhizomania).

A compound of formula (I) may move acropetally, basipetally or locally in plant
10 tissue to be active against one or more fungi.

The invention therefore provides a method of combating or controlling phytopathogenic fungi which comprises applying a fungicidally effective amount of a compound of formula (I), or a composition containing a compound of formula (I), to a plant, to a seed of a plant, to the locus of the plant or seed or to soil or any other plant growth medium, e.g.

15 nutrient solution.

The term "plant" as used herein includes seedlings, bushes and trees. Furthermore, the fungicidal method of the invention includes protectant, curative, systemic, eradicator and antispore treatments.

The compound of formula (I) is preferably used for agricultural, horticultural and
20 turfgrass purposes in the form of a composition.

In order to apply a compound of formula (I) to a plant, to a seed of a plant, to the locus of the plant or seed or to soil or any other growth medium, a compound of formula (I) is usually formulated into a composition which includes, in addition to the compound of formula (I), a suitable inert diluent or carrier and, optionally, a surface active agent (SFA).
25 SFAs are chemicals which are able to modify the properties of an interface (for example, liquid/solid, liquid/air or liquid/liquid interfaces) by lowering the interfacial tension and thereby leading to changes in other properties (for example dispersion, emulsification and wetting). It is preferred that all compositions (both solid and liquid formulations) comprise, by weight, 0.0001 to 95%, more preferably 1 to 85%, for example 5 to 60%, of a compound
30 of formula (I). The composition is generally used for the control of fungi such that a compound of formula (I) is applied at a rate of from 0.1g to 10kg per hectare, preferably from 1g to 6kg per hectare, more preferably from 1g to 1kg per hectare.

When used in a seed dressing, a compound of formula (I) is used at a rate of 0.0001g to 10g (for example 0.001g or 0.05g), preferably 0.005g to 10g, more preferably 0.005g to 4g, per kilogram of seed.

5 In another aspect the present invention provides a biocidal composition comprising a biocidally effective amount of a compound of formula (I) and a suitable carrier or diluent therefor.

In a preferred embodiment of this aspect the present invention provides an antiviral composition comprising an antivirally effective amount of a compound of formula (I) and a suitable carrier or diluent therefor.

10 In a further preferred embodiment of this aspect the present invention provides an anti-cancer composition comprising an effective amount of a compound of formula (I) which is effective in reducing the proliferation of cancer cells and a suitable carrier or diluent therefor.

15 In a particularly preferred embodiment of this aspect the present invention provides a fungicidal composition comprising a fungicidally effective amount of a compound of formula (I) and a suitable carrier or diluent therefor.

In a still further aspect the invention provides a method of combating and controlling viruses which comprises treating the viruses or the locus of the viruses with an antivirally effective amount of a composition comprising a compound of formula (I).

20 In a still further aspect the invention provides a method of combating and controlling cancer cells which comprises treating the cancer cells or the locus of the cancer cells with an effective amount of a composition comprising a compound of formula (I) said amount being effective in reducing the proliferation of said cancer cells.

25 In a still further preferred aspect the invention provides a method of combating and controlling fungi which comprises treating the fungi or the locus of the fungi with a fungicidally effective amount of a composition comprising a compound of formula (I).

The compositions can be chosen from a number of formulation types, including dustable powders (DP), soluble powders (SP), water soluble granules (SG), water dispersible granules (WG), wettable powders (WP), granules (GR) (slow or fast release), soluble
30 concentrates (SL), oil miscible liquids (OL), ultra low volume liquids (UL), emulsifiable concentrates (EC), dispersible concentrates (DC), emulsions (both oil in water (EW) and water in oil (EO)), micro-emulsions (ME), suspension concentrates (SC), aerosols, fogging/smoke formulations, capsule suspensions (CS) and seed treatment formulations.

The formulation type chosen in any instance will depend upon the particular purpose envisaged and the physical, chemical and biological properties of the compound of formula (I).

Dustable powders (DP) may be prepared by mixing a compound of formula (I) with one or more solid diluents (for example natural clays, kaolin, pyrophyllite, bentonite, alumina, montmorillonite, kieselguhr, chalk, diatomaceous earths, calcium phosphates, calcium and magnesium carbonates, sulphur, lime, flours, talc and other organic and inorganic solid carriers) and mechanically grinding the mixture to a fine powder.

Soluble powders (SP) may be prepared by mixing a compound of formula (I) with one or more water-soluble inorganic salts (such as sodium bicarbonate, sodium carbonate or magnesium sulphate) or one or more water-soluble organic solids (such as a polysaccharide) and, optionally, one or more wetting agents, one or more dispersing agents or a mixture of said agents to improve water dispersibility/solubility. The mixture is then ground to a fine powder. Similar compositions may also be granulated to form water soluble granules (SG).

Wettable powders (WP) may be prepared by mixing a compound of formula (I) with one or more solid diluents or carriers, one or more wetting agents and, preferably, one or more dispersing agents and, optionally, one or more suspending agents to facilitate the dispersion in liquids. The mixture is then ground to a fine powder. Similar compositions may also be granulated to form water dispersible granules (WG).

Granules (GR) may be formed either by granulating a mixture of a compound of formula (I) and one or more powdered solid diluents or carriers, or from pre-formed blank granules by absorbing a compound of formula (I) (or a solution thereof, in a suitable agent) in a porous granular material (such as pumice, attapulgite clays, fuller's earth, kieselguhr, diatomaceous earths or ground corn cobs) or by adsorbing a compound of formula (I) (or a solution thereof, in a suitable agent) on to a hard core material (such as sands, silicates, mineral carbonates, sulphates or phosphates) and drying if necessary. Agents which are commonly used to aid absorption or adsorption include solvents (such as aliphatic and aromatic petroleum solvents, alcohols, ethers, ketones and esters) and sticking agents (such as polyvinyl acetates, polyvinyl alcohols, dextrans, sugars and vegetable oils). One or more other additives may also be included in granules (for example an emulsifying agent, wetting agent or dispersing agent).

Dispersible Concentrates (DC) may be prepared by dissolving a compound of formula (I) in water or an organic solvent, such as a ketone, alcohol or glycol ether. These

solutions may contain a surface active agent (for example to improve water dilution or prevent crystallisation in a spray tank).

Emulsifiable concentrates (EC) or oil-in-water emulsions (EW) may be prepared by dissolving a compound of formula (I) in an organic solvent (optionally containing one or more wetting agents, one or more emulsifying agents or a mixture of said agents). Suitable organic solvents for use in ECs include aromatic hydrocarbons (such as alkylbenzenes or alkylnaphthalenes, exemplified by SOLVESSO 100, SOLVESSO 150 and SOLVESSO 200; SOLVESSO is a Registered Trade Mark), ketones (such as cyclohexanone or methylcyclohexanone) and alcohols (such as benzyl alcohol, furfuryl alcohol or butanol), N-alkylpyrrolidones (such as N-methylpyrrolidone or N-octylpyrrolidone), dimethyl amides of fatty acids (such as C₈-C₁₀ fatty acid dimethylamide) and chlorinated hydrocarbons. An EC product may spontaneously emulsify on addition to water, to produce an emulsion with sufficient stability to allow spray application through appropriate equipment. Preparation of an EW involves obtaining a compound of formula (I) either as a liquid (if it is not a liquid at room temperature, it may be melted at a reasonable temperature, typically below 70°C) or in solution (by dissolving it in an appropriate solvent) and then emulsifying the resultant liquid or solution into water containing one or more SFAs, under high shear, to produce an emulsion. Suitable solvents for use in EWs include vegetable oils, chlorinated hydrocarbons (such as chlorobenzenes), aromatic solvents (such as alkylbenzenes or alkylnaphthalenes) and other appropriate organic solvents which have a low solubility in water.

Microemulsions (ME) may be prepared by mixing water with a blend of one or more solvents with one or more SFAs, to produce spontaneously a thermodynamically stable isotropic liquid formulation. A compound of formula (I) is present initially in either the water or the solvent/SFA blend. Suitable solvents for use in MEs include those hereinbefore described for use in ECs or in EWs. An ME may be either an oil-in-water or a water-in-oil system (which system is present may be determined by conductivity measurements) and may be suitable for mixing water-soluble and oil-soluble pesticides in the same formulation. An ME is suitable for dilution into water, either remaining as a microemulsion or forming a conventional oil-in-water emulsion.

Suspension concentrates (SC) may comprise aqueous or non-aqueous suspensions of finely divided insoluble solid particles of a compound of formula (I). SCs may be prepared by ball or bead milling the solid compound of formula (I) in a suitable medium, optionally with one or more dispersing agents, to produce a fine particle suspension of the compound.

One or more wetting agents may be included in the composition and a suspending agent may be included to reduce the rate at which the particles settle. Alternatively, a compound of formula (I) may be dry milled and added to water, containing agents hereinbefore described, to produce the desired end product.

5 Aerosol formulations comprise a compound of formula (I) and a suitable propellant (for example *n*-butane). A compound of formula (I) may also be dissolved or dispersed in a suitable medium (for example water or a water miscible liquid, such as *n*-propanol) to provide compositions for use in non-pressurised, hand-actuated spray pumps.

10 A compound of formula (I) may be mixed in the dry state with a pyrotechnic mixture to form a composition suitable for generating, in an enclosed space, a smoke containing the compound.

15 Capsule suspensions (CS) may be prepared in a manner similar to the preparation of EW formulations but with an additional polymerisation stage such that an aqueous dispersion of oil droplets is obtained, in which each oil droplet is encapsulated by a polymeric shell and contains a compound of formula (I) and, optionally, a carrier or diluent therefor. The polymeric shell may be produced by either an interfacial polycondensation reaction or by a coacervation procedure. The compositions may provide for controlled release of the compound of formula (I) and they may be used for seed treatment. A compound of formula (I) may also be formulated in a biodegradable polymeric matrix to provide a slow, controlled
20 release of the compound.

25 A composition may include one or more additives to improve the biological performance of the composition (for example by improving wetting, retention or distribution on surfaces; resistance to rain on treated surfaces; or uptake or mobility of a compound of formula (I)). Such additives include surface active agents, spray additives based on oils, for example certain mineral oils or natural plant oils (such as soy bean and rape seed oil), and blends of these with other bio-enhancing adjuvants (ingredients which may aid or modify the
30 action of a compound of formula (I)).

 A compound of formula (I) may also be formulated for use as a seed treatment, for example as a powder composition, including a powder for dry seed treatment (DS), a water soluble powder (SS) or a water dispersible powder for slurry treatment (WS), or as a liquid
30 composition, including a flowable concentrate (FS), a solution (LS) or a capsule suspension (CS). The preparations of DS, SS, WS, FS and LS compositions are very similar to those of, respectively, DP, SP, WP, SC and DC compositions described above. Compositions for

treating seed may include an agent for assisting the adhesion of the composition to the seed (for example a mineral oil or a film-forming barrier).

Wetting agents, dispersing agents and emulsifying agents may be SFAs of the cationic, anionic, amphoteric or non-ionic type.

5 Suitable SFAs of the cationic type include quaternary ammonium compounds (for example cetyltrimethyl ammonium bromide), imidazolines and amine salts.

 Suitable anionic SFAs include alkali metals salts of fatty acids, salts of aliphatic monoesters of sulphuric acid (for example sodium lauryl sulphate), salts of sulphonated aromatic compounds (for example sodium dodecylbenzenesulphonate, calcium
10 dodecylbenzenesulphonate, butylnaphthalene sulphonate and mixtures of sodium di-*isopropyl*- and tri-*isopropyl*-naphthalene sulphonates), ether sulphates, alcohol ether sulphates (for example sodium laureth-3-sulphate), ether carboxylates (for example sodium laureth-3-carboxylate), phosphate esters (products from the reaction between one or more fatty alcohols and phosphoric acid (predominately mono-esters) or phosphorus pentoxide
15 (predominately di-esters), for example the reaction between lauryl alcohol and tetraphosphoric acid; additionally these products may be ethoxylated), sulphosuccinamates, paraffin or olefine sulphonates, taurates and lignosulphonates.

 Suitable SFAs of the amphoteric type include betaines, propionates and glycines.

 Suitable SFAs of the non-ionic type include condensation products of alkylene
20 oxides, such as ethylene oxide, propylene oxide, butylene oxide or mixtures thereof, with fatty alcohols (such as oleyl alcohol or cetyl alcohol) or with alkylphenols (such as octylphenol, nonylphenol or octylcresol); partial esters derived from long chain fatty acids or hexitol anhydrides; condensation products of said partial esters with ethylene oxide; block polymers (comprising ethylene oxide and propylene oxide); alkanolamides; simple esters (for
25 example fatty acid polyethylene glycol esters); amine oxides (for example lauryl dimethyl amine oxide); and lecithins.

 Suitable suspending agents include hydrophilic colloids (such as polysaccharides, polyvinylpyrrolidone or sodium carboxymethylcellulose) and swelling clays (such as bentonite or attapulgate).

30 A compound of formula (I) may be applied by any of the known means of applying fungicidal compounds. For example, it may be applied, formulated or unformulated, to any part of the plant, including the foliage, stems, branches or roots, to the seed before it is planted or to other media in which plants are growing or are to be planted (such as soil

surrounding the roots, the soil generally, paddy water or hydroponic culture systems), directly or it may be sprayed on, dusted on, applied by dipping, applied as a cream or paste formulation, applied as a vapour or applied through distribution or incorporation of a composition (such as a granular composition or a composition packed in a water-soluble bag) in soil or an aqueous environment.

A compound of formula (I) may also be injected into plants or sprayed onto vegetation using electrodynamic spraying techniques or other low volume methods, or applied by land or aerial irrigation systems.

Compositions for use as aqueous preparations (aqueous solutions or dispersions) are generally supplied in the form of a concentrate containing a high proportion of the active ingredient, the concentrate being added to water before use. These concentrates, which may include DCs, SCs, ECs, EWs, MEs SGs, SPs, WPs, WGs and CSs, are often required to withstand storage for prolonged periods and, after such storage, to be capable of addition to water to form aqueous preparations which remain homogeneous for a sufficient time to enable them to be applied by conventional spray equipment. Such aqueous preparations may contain varying amounts of a compound of formula (I) (for example 0.0001 to 10%, by weight) depending upon the purpose for which they are to be used.

A compound of formula (I) may be used in mixtures with fertilisers (for example nitrogen-, potassium- or phosphorus-containing fertilisers). Suitable formulation types include granules of fertiliser. The mixtures suitably contain up to 25% by weight of the compound of formula (I).

The invention therefore also provides a fertiliser composition comprising a fertiliser and a compound of formula (I).

The compositions of this invention may contain other compounds having biological activity, for example micronutrients or compounds having similar or complementary fungicidal activity or which possess plant growth regulating, herbicidal, insecticidal, nematocidal or acaricidal activity.


By including another fungicide, the resulting composition may have a broader spectrum of activity or a greater level of intrinsic activity than the compound of formula (I) alone. Further the other fungicide may have a synergistic effect on the fungicidal activity of the compound of formula (I).

The compound of formula (I) may be the sole active ingredient of the composition or it may be admixed with one or more additional active ingredients such as a pesticide,

fungicide, synergist, herbicide or plant growth regulator where appropriate. An additional active ingredient may: provide a composition having a broader spectrum of activity or increased persistence at a locus; synergise the activity or complement the activity (for example by increasing the speed of effect or overcoming repellency) of the compound of formula (I); or help to overcome or prevent the development of resistance to individual components. The particular additional active ingredient will depend upon the intended utility of the composition.

Examples of fungicidal compounds which may be included in the composition of the invention are AC 382042 (*N*-(1-cyano-1,2-dimethylpropyl)-2-(2,4-dichlorophenoxy) propionamide), acibenzolar-S-methyl, alanycarb, aldimorph, anilazine, azaconazole, azafenidin, azoxystrobin, benalaxyl, benomyl, biloxazol, bitertanol, blasticidin S, bromuconazole, bupirimate, captafol, captan, carbendazim, carbendazim chlorhydrate, carboxin, carpropamid, carvone, CGA 41396, CGA 41397, chinomethionate, chlorbenzthiazole, chlorothalonil, chlorozolinate, clozylacon, copper containing compounds such as copper oxychloride, copper oxyquinolate, copper sulphate, copper tallate, and Bordeaux mixture, cyamidazosulfamid, cyflufenamid, cymoxanil, cyproconazole, cyprodinil, debacarb, di-2-pyridyl disulphide 1,1'-dioxide, dichlofluanid, diclocymet, diclomezine, dicloran, diethofencarb, difenoconazole, difenzoquat, diflumetorim, *O,O*-di-*iso*-propyl-S-benzyl thiophosphate, dimefluazole, dimetconazole, dimethirimol, dimethomorph, dimoxystrobin, diniconazole, dinocap, dithianon, dodecyl dimethyl ammonium chloride, dodemorph, dodine, doguadine, edifenphos, epoxiconazole, ethaboxam, ethirimol, ethyl (*Z*)-*N*-benzyl-*N*[(methyl(methyl-thioethylideneaminooxycarbonyl)amino)thio]- β -alaninate, etridiazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fenfuram, fenhexamid, fenpiclonil, fenpropidin, fenpropimorph, fentin acetate, fentin hydroxide, ferbam, ferimzone, fluazinam, fludioxonil, flumetover, flumorph, fluoroimide, fluoxastrobin, fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpet, fosetyl-aluminium, fuberidazole, furalaxyl, furametpyr, guazatine, hexaconazole, hydroxyisoxazole, hymexazole, imazalil, imibenconazole, iminoctadine, iminoctadine triacetate, ipconazole, iprobenfos, iprodione, iprovalicarb, isopropanyl butyl carbamate, isoprothiolane, kasugamycin, kresoxim-methyl, LY186054, LY211795, LY 248908, mancozeb, maneb, mefenoxam, mepanipyrim, mepronil, metalaxyl, metconazole, metiram, metiram-zinc, metominostrobin, metrafenone, MON65500 (*N*-allyl-4,5-dimethyl-2-trimethylsilylthiophene-3-carboxamide), myclobutanil, NTN0301, neoasozin, nickel dimethyldithiocarbamate, nicobifen, nitrothale-isopropyl, nuarimol,

ofurace, organomercury compounds, oxadixyl, oxasulfuron, oxolinic acid, oxpoconazole, oxycarboxin, pefurazoate, penconazole, pencycuron, phenazin oxide, phosphorus acids, phthalide, picoxystrobin, polyoxin D, polyram, probenazole, prochloraz, procymidone, propamocarb, propamocarb hydrochloride, propiconazole, propineb, propionic acid, pyraclostrobin, pyrazophos, pyrifenox, pyrimethanil, pyroquilon, pyroxyfur, pyrrolnitrin, quaternary ammonium compounds, quinomethionate, quinoxifen, quintozone, sipconazole, sodium pentachlorophenate, spiroxamine, streptomycin, sulphur, tebuconazole, tecloftalam, tecnazene, tetraconazole, thiabendazole, thifluzamide, 2-(thiocyanomethylthio)-benzothiazole, thiophanate-methyl, thiram, tiadinil, timibenconazole, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, triazbutil, triazoxide, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, validamycin A, vapam, vinclozolin, XRD-563, zineb, ziram, zoxamide and compounds of the formulae:



The compound of formula (I) may be mixed with soil, peat or other rooting media for the protection of plants against seed-borne, soil-borne or foliar fungal diseases.

Some mixtures may comprise active ingredients which have significantly different physical, chemical or biological properties such that they do not easily lend themselves to the same conventional formulation type. In these circumstances other formulation types may be prepared. For example, where one active ingredient is a water insoluble solid and the other a water insoluble liquid, it may nevertheless be possible to disperse each active ingredient in the same continuous aqueous phase by dispersing the solid active ingredient as a suspension (using a preparation analogous to that of an SC) but dispersing the liquid active ingredient as an emulsion (using a preparation analogous to that of an EW). The resultant composition is a suspoemulsion (SE) formulation.

The compound of the invention may be formulated for administration in any convenient way for use in veterinary or human medicine and the invention therefore includes within its scope pharmaceutical compositions comprising a compound of formula (I) adapted for use in veterinary or human medicine. Such compositions may be presented for use in conventional manner with the aid of one or more suitable pharmaceutically or veterinary acceptable carriers or excipients.

The compositions of the invention include those in a form formulated for parenteral, oral, rectal, topical or implant use.

For use in agriculture, horticulture or veterinary medicine it may be desirable to use the whole fermentation broth as a source of the active compound of the invention without separation into the individual compound of formula (I). It may be suitable to use dried broth containing mycelia or to use lysed mycelia, live or dead mycelia separated from the broth using solid/liquid separation or evaporation techniques or to use the fermentation broth remaining after separation of the mycelia. If desired the broth or mycelia may be formulated into compositions including conventional inert carriers, excipients or diluents.

It will be understood to the man skilled in the art that in general the compound of the invention may be used to combat infections by applying to the organism responsible for the infection or a location thereof an effective amount of a compound of formula (I).

In a further aspect the invention provides a process for the production of a compound of formula (I) which comprises the step of cultivating an organism of the strain *Streptomyces malaysiensis* capable of producing a compound of formula (I) whereby said compound is produced, and if desired isolating said compound therefrom.

Taxonomic studies have shown that a particular microorganism capable of producing a compound of formula (I) is a new strain of the microorganism *Streptomyces malaysiensis* hereinafter referred to as *Streptomyces malaysiensis* JHCC 553434. A sample of this microorganism, which is a soil isolate, was deposited on 17 December 2001 at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Mascheroder Weg 1b, D-38124 Braunschweig, Germany and has been assigned the Accession Number DSM 14702. The strain has been deposited under the terms and conditions of the Budapest Treaty and has been placed under restricted access under the terms of Rule 28(4) of the European Patent Convention. This new strain of *Streptomyces malaysiensis*, *Streptomyces malaysiensis* JHCC 553434 (DSM 14702) provides a further feature of this invention. The invention also extends to further new strains of *Streptomyces malaysiensis* which possess the same essential morphological and cultural characteristics as *Streptomyces malaysiensis* JHCC 553434 (DSM 14702) and which are capable of producing a compound of formula (I).

The essential morphological and cultural characteristics assigned to *Streptomyces malaysiensis* JHCC 553434 (DSM 14702) are listed below:

- Diagnostic aminoacid of peptidoglycan is: LL-diaminopimeli(ni)c acid
- Mycolic acids are absent

- Iso- and ante-iso fatty acids found, but no 10-methyl branched fatty acids
- Base-sequence of 16s rDNA: similarity to other sequences of *Streptomyces* species >95%
- Colour of colony: grey to dark-brown aerial mycelium and beige dirty-yellow substrate mycelium
- Formation of melanin or other diagnostic pigments not found
- Starch is hydrolysed
- Morphology: spores in short and narrow coils
- Physiology: according to Kaempfer et al. (J. Gen Microbiol. **13** 1831-1891 (1991)): utilisation of carbon sources => 100% cluster 18 : *S. violaceusniger*
- Fatty acid -pattern: enough similarity (0.675) to *S. violaceusniger*
- 16s rDNA sequencing: comparison of partial sequence to *Streptomyces* library gave a similarity of 100% to *Streptomyces malaysiensis*, a member of group *S. violaceusniger*
- Riboprint-pattern: enough similarity to group *S. violaceusniger*. Species *S. hygroscopicus* 0.83 and *S. malyasiensis* 0.82.

The species *Streptomyces malaysiensis* was recently described in literature: International Journal of Systematic Bacteriology (1999), 49, 1395-1402 by Amira Al-Tai et al.

The invention extends also to mutant strains of *Streptomyces malaysiensis* JHCC 553434 (DSM 14702) capable of producing a compound of formula (I) . Mutants may be naturally occurring or may be produced by mutagenesis of the parent strain using conventional mutagenesis techniques such as genetic techniques such as recombination, transformation and selective techniques for spontaneous mutants; ionising radiation; chemical methods; heat;.

The production of a compound of formula (I) by cultivation of a suitable strain of *Streptomyces malaysiensis* may be effected by conventional means i.e. by culturing the microorganism in the presence of assimilable sources of carbon, nitrogen and mineral salts.

The strain of *Streptomyces malaysiensis* according to the invention is preferably *Streptomyces malaysiensis* JHCC 553434 (DSM 14702).

Assimilable sources of nitrogen, carbon, and minerals may be provided by either simple or complex nutrients. Sources of carbon will generally include amino acids; alkanes; molasses; glycerides; glycerol; glucose, maltose, lactose, sucrose, fructose, starch, dextrin,

carboxylic acids, alcohols, and vegetable oils. Sources of carbon will generally comprise from 0.5% to 10% by weight of the fermentation medium.

Sources of nitrogen will generally include soya bean meal, corn steep liquors, yeast extracts, cottonseed meal, peptones, malt extract, molasses, casein, amino acid mixture, ammonia, ammonium salts or nitrates, urea and other amides. Sources of nitrogen will generally comprise from 0.1% to 10% by weight of the fermentation medium.

Nutrient mineral salts which may be incorporated into the culture medium include for example, salts capable of generating sodium, potassium, ammonium, iron, magnesium, zinc, nickel, cobalt, manganese, vanadium, chromium, calcium, copper, molybdenum, boron, phosphate, sulphate, chloride and carbonate ions.

Cultivation will desirably take place with aeration and agitation. The cultivation will generally be carried out in the pH range 5 to 8, preferably 6 to 7, most preferably pH 6.5. The microorganism of the invention will generally be cultivated at a temperature of from 20° to 50°C, preferably from 25° to 40°C and most especially at 28° or 35°C. The fermentation may be carried out for a period of 2- 14 days, e.g. 6 to 8 days.

Where it is desired to separate the compound of the invention from the whole fermentation broth this may be achieved using conventional isolation and separation techniques. The compound of the invention is found predominantly in the fermentation broth but small amounts may be found in the mycelia of the cells.

In a further aspect the invention provides a compound of formula (I) in the form of a whole fermentation broth containing such a compound; the solids of a whole fermentation broth containing such a compound; intact or lysed mycelia separated from a whole fermentation broth containing such a compound; or the solids of such a whole fermentation broth containing such a compound after separation of intact or lysed mycelia; or such a whole fermentation broth containing such a compound after the separation of the mycelia and all solids.

In a yet further aspect the invention provides a composition comprising an effective amount of a compound of formula (I) which may be in a form according to the above aspect of the invention.

The compound of the invention may be isolated and separated by a variety of fractionation techniques, for example, adsorption-elution, precipitation, fractional crystallisation and solvent extraction which may be combined in various ways.

Centrifugation and chromatography have been found to be most suitable for isolating and separating the compound of the invention.

Following the fermentation, the mycelia may be harvested using conventional techniques such as filtration or centrifugation. The compound of the invention is found
5 mainly in the aqueous medium and may be separated from the mycelia, for example by centrifugation, followed by extraction of the mycelium with water or an appropriate water miscible solvent such as a lower ketone e.g. acetone; a lower alcohol e.g. methanol or ethanol; or a lower diol.

Generally more than one extraction is desirable to achieve optimum recovery.

10 Preferably the first extraction is performed using water or a water soluble solvent, preferably methanol. The compound of the invention may then be recovered as a crude extract by removal of the solvent e.g. by evaporation. The concentrate may then be redissolved in an aqueous medium e.g. water, and further purified by further centrifugation of any residual solids. Removal of the water or water miscible solvent yields a compound of formula (I).

15 Purification of the compound of the invention may be effected by conventional techniques such as for example chromatography, preferably high performance liquid chromatography, on a suitable support. The support may be in the form of a bed or preferably packed in a column.

20 A solution of the compound in a suitable solvent will generally be loaded on to the silica, if desired after reducing the volume of the solvent. The column may optionally be washed and eluted with a solvent of suitable polarity. In the case of silica, alcohols such as methanol in combination with water may be used.

25 Elution and separation of the compound of the invention may be monitored by conventional techniques such as chromatography preferably high performance liquid chromatography.

The compound of the invention may be further purified by crystallisation and the invention extends also to the compound of formula (I) in crystalline form.

The invention further extends to a process for combating fungi or viruses or cancer cells comprising exposing said viruses or fungi or cancer cells to a compound of formula (I).

30 The invention extends also to the use of a compound of formula I with a biological system which has modified or enhanced resistance to fungi and/or viruses. The biological system may for example be a plant, or may be a mammalian system such as an organ, tissue or cell thereof. In the case of a plant system the enhanced resistance may have been

introduced into the plant using conventional breeding techniques where plants showing enhanced resistance are selected over many generations. Alternatively or as well as, the enhanced resistance may be due to the introduction into said plant by genetic modification recombinant DNA techniques of one or more DNA sequences, the expression of which enhances the resistance of the host plant. This approach is often referred to as integrated crop management.

In a further aspect therefore the invention provides a method of controlling fungal infection of a plant or part thereof said plant having been genetically modified to enhance resistance to fungi either by selective breeding and/or preferably by genetic modification where one or more DNA sequences, the expression of which enhances the resistance of the plant to fungi have been introduced into said plant using recombinant DNA techniques, said method comprising exposing said plant or part thereof to a compound of formula(I).

In a further aspect therefore the invention provides a method of controlling viral infection of a plant or part thereof said plant having been genetically modified to enhance resistance to viruses either by selective breeding and/or preferably by genetic modification where one or more DNA sequences, the expression of which enhances the resistance of the plant to viruses have been introduced into said plant using recombinant DNA techniques, said method comprising exposing said plant or part thereof to a compound of formula(I).

In a yet further aspect the invention provides the use of a compound of formula (I) as a biocidal agent.

In a preferred embodiment of this aspect the invention provides the use of a compound of formula (I) as an anti-fungal agent.

In a further embodiment of this aspect the invention provides the use of a compound of formula (I) as an anti-viral agent.

In a further embodiment of this aspect the invention provides the use of a compound of formula (I) as an anti-cancer agent.

The invention is illustrated by the following Examples:

Example 1

Standard Procedure for Production of Compound of Formula (I)

Shake flasks in M3 media pH controlled (MES)

Producer strain *Streptomyces malaysiensis*. JHCC-553434 was grown on agar slants and plates. It was transferred to TSM-40 and TSM-7/2 to grow up for strain preservation. Better

growth was found in TSM-7/2 and it was used for routine strain preservation procedure. A stock of -80°C cryovials was produced as start material for all following experiments. HPLC-MS trials using the supernatant of those cultures (4-6 days at 35°C) show production of a compound of formula (I).

5 **TSM-40**

for 1l

Glucose (Merck) 4g

Malz Extract (Difco) 10g

Yeast Extract 4g

10 pH corrected before sterilisation to 7

TSM-7/2

for 1l

Mannit 20g

15 Soytone 10g

Sojabeens Oil 2.5g

K₂HPO₄ 0.35g

CaCO₃ 0.5g

pH corrected before sterilisation to 7

20

Setup of production in shake flasks

Inoculums: As starting material good grown 1 x ISP-2 agar plates were used and incubated at 35°C about 5 days.

25 **Seed culture**

VK1: 1 x SC-02 medium (100 ml in 1 baffled EMK 500 ml) was inoculated with biomass from agar plates and incubated for 3 days at 35°C and 120 rpm on a rotary shaker.

30 **Main culture:** 150 x M3 medium (60 ml in 1 baffled EMK 500 ml) were inoculated with 5 vol% of SC-01 and incubated on the rotary shaker at 35°C and 120 rpm. Media were buffered using 100 mM MES buffer (Good's buffer) and pH was

adjusted to 6.5 (NaOH) prior to autoclaving. (MES Buffer is 2-N-Morpholino ethanesulfonic acid Hydrate , minimum 99.5 % titration .
C₆H₁₃NO₄S 1M = 195.2g. pKa = 6.1 at 25°C, pH range 5.5-6.7 Producer
SIGMA , ART. Nr. M8250)

- 5 Samples and Analysis: Samples were checked on days: 4, 5 and 6. Growth was recorded and
1 ml of samples were centrifuged (Biofuge A- Heraeus Sepatec, 10 min 10
000 rpm, RT). Supernatant was separated and biomass was discarded. All
amount of supernatants were filtrated. Clear supernatants were transferred to
HPLC vials and measured in a Waters LC/MS system (HPLC: Alliance 2790
10 with photodiode array detector, MS: micromass ZQ).

Harvest: and Analysis: After 6 days all flasks were harvested and centrifuged. Supernatant
and Biomass were stored at -20 C.

15 ***MEDIA:***

SC-02

For 1l medium:

	Glucose	(Merck)	10 g
	Starch soluble	(Merck)	10 g
20	Peptone	(Sigma)	10 g
	Yeast extract	(Difco)	2 g
	MgSO ₄ x 7H ₂ O	(Merck)	1 g
	pH 6.5		

25 **M3**

For 1 l medium:

	Glucose	20g
	Starch sol.	10g
	Sojaflower	25g
30	Yeast Extract	4g

Meat Extract 1g
Component A 10 ml

Component A:

5 For 1l add before sterilization:

NaCl	0.5 g
CaCO ₃	0.5 g
MgSO ₄ x 7H ₂ O	0.5 g
KH ₂ PO ₄	50 mg
10 FeCl ₃ x 6H ₂ O	16.7 mg
ZnSO ₄ x 7H ₂ O	21.2 mg
CaCl ₂ x 2H ₂ O	6.5 mg
CuSO ₄ x 5H ₂ O	5 mg
MnCl ₂ x 4 H ₂ O	5 mg

15


In general 1l of 100x concentrated stock solution is prepared. From stock solution 10 ml /l is used. Stock solution has to be stored at 4°C.

Example 2

20

Isolation and physical-/chemical characterisation of Compound of Formula (I)

23g of lyophilisate, which corresponds to 1 l of fermentation broth of *Streptomyces malaysiensis* JHCC-553434 prepared as described in Example 1, are dissolved in water and transferred on a G-25 column (Sephadex, 70mm x 780 mm, 1 ml/min). After elution with
25 water, the fraction containing the compound of formula (I) is lyophilised. The residue, 0.678g, is purified on a medium-pressure chromatography on silica gel (Silica 32-63, 60 Angström, 55 mm x 500 mm, 22 ml/min) with a gradient of acetonitrile:water and yields after evaporation, 0.143g of enriched material. Further separation by means of high-pressure chromatography on RP silica gel (Kromasil 100, C18, 10 µm, 20 mm x 250 mm, 10 ml/min),
30 with a gradient of acetonitrile:water, yields 0.034g of pure compound of formula (I).

Example 3**Characterising Structural Data for Compound of Formula (I)****5 NMR Data**

^1H spectrum, run 2% w/v solution in D_2O , referenced to residual HOD peak at δ 4.63. Data in { } was run as 0.5% w/v solution in $\text{D}_6\text{-DMSO}$, referenced to HD5-DMSO at δ 2.49. Spectra were run on a JEOL GX400 Spectrometer operating at 399.6 MHz

10

δ	Carbon to which H's bonded	Multiplicity and Number of H's	Coupling constant in Hz
3.26	C8	s (3H)	
3.37	C7	t (1H)	11.4
3.48	C3	dd (1H)	10.8, 3.8
3.67	C6	ddd (1H)	11.5, 5.6, 4.3
3.83	C4	dd (1H)	10.8, 3.8
3.92	C7	dd (1H)	11.4, 4.8
4.14	C2	d {d}(1H)	4.8, {4.0}
4.71	C1	dd (1H)	1.3, 0.7
4.80	C5	m (1H)	
{5.12}	C2	d (1H) OH	4.0
{6.30}	C9	d (1H) NH	10.5
7.41	Uracil C4	d (1H)	1.3
{10.85}		Uracil NH's and urea NH ₂	

s = singlet

- d = doublet
 dd = double doublet
 ddd = double double doublet
 t = triplet
 5 m = multiplet


- 10 ^{13}C spectrum, run 2% w/v solution in D_2O , referenced to D6-acetone $\{(\text{CD}_3)_2\text{CO}\}$ at δ 29.8, using JEOL GX400 NMR spectrometer operating at 100.4 MHz

δ	Carbon	Multiplicity
47.3	C5	d (broadened)
56.2	C8	q
65.2	C7	t
70.8	C2	d
73.2	C3	d
73.3	C6	d
74.4	C4	d
83.4	C1	q
111.4	Uracil C5	s
139.0	Uracil C4	d
152.6	Uracil C2	s
161.2	C9	s
164.5	Uracil C6	s

- s = singlet
 15 d = doublet
 t = triplet
 q = quartet

Terms above relate to signals off-resonance multiplicity as determined by DEPT (135) spectrum.

On the basis of the above nmr data we believe the stereochemistry to be as shown below



Mass Spectrometry

Ionisation by fast atom bombardment in glycerol and water gives an m/z of 343 $[M+H]^+$ and
10 m/z of 365 $[M+Na]^+$, consistent with a molecular weight of 342.

Accurate mass measurement gave a molecular weight as 342.1197 mass units consistent with the molecular formula of $C_{13}H_{18}O_7N_4$.

15 EXAMPLE 4

This Example illustrates the fungicidal properties of a compound of formula (I). The compound was tested against a variety of foliar fungal diseases of plants. The technique employed was as follows.

Plants were either grown in John Innes Potting Compost (No.1 or 2) in 4cm diameter,
20 3.5cm depth minipots or on an artificial, cellulose based growing medium. The test compound was individually prepared as a solution in acetone. This was diluted in reverse osmosis water to a final concentration of 100ppm in a 1:2 mixture of acetone:water (that is, 1mg of compound in a final volume of 10ml) immediately before use. TWEEN 20 (at a final concentration of 0.05% by volume) was added with the water to improve retention of the
25 spray deposit. TWEEN is a registered trade mark.

A compound of formula (I) was applied as a foliar application where the chemical solution was applied to the foliage of the test plants by spraying the plant to maximum droplet retention.

These tests were carried out against *Phytophthora infestans lycopersici* (PHYTIN) on tomatoes; *Blumeria graminis f.sp. tritici* (ERYSGT), and *Puccinia triticina* (PUCCRT) on wheat. Each treatment was applied to two replicate plants for *Phytophthora infestans lycopersici*. In tests on *Blumeria graminis f.sp. tritici*, and *Puccinia triticina* two replicates, each containing 3 plants were used for each treatment. The plants were inoculated with either a calibrated fungal spore suspension or a "dusting" with dry spores 6 hours or one day after chemical application.

After chemical application and inoculation, the plants were incubated under high humidity conditions (except those inoculated with *Blumeria graminis f.sp. tritici*) and then put into an appropriate environment to allow infection to proceed, until the disease was ready for assessment. The time period between chemical application and assessment varied from six to nine days according to the disease and environment. However, each individual disease was assessed after the same time period for all compounds.

Assessments were performed on each of two leaves on each of the replicate plants for *Phytophthora infestans lycopersici*. For *Blumeria graminis f.sp. tritici* and *Puccinia triticina* assessments were carried out collectively on the plants in each replicate and averaged to give one result per replicate.

The disease level present (the percentage leaf area covered by actively sporulating disease) was assessed visually. For each treatment, the assessed values for all its replicates were meaned to provide mean disease values. Untreated control plants were assessed in the same manner. The data were then processed (see formula below) to calculate a PRCO (Percentage Disease Reduction from Control) value.

BANDED ASSESSMENT METHOD AND CALCULATION OF PRCO VALUES

The mean disease values are banded in the manner shown below. If the disease level value falls exactly mid-way between two of the points, the result will be the lower of the two points.

0 =	0% disease present	10 =	5.1-10% disease present
1 =	0.1-1% disease present	20 =	10.1-20% disease present
3 =	1.1-3% disease present	30 =	20.1-30% disease present
5 =	3.1-5% disease present	60 =	30.1-60% disease present
		90 =	60.1-100% disease present

An example of a typical banded calculation is as follows:

Mean disease level for treatment A = 25%

Therefore banded mean disease level for treatment A = 30

Mean disease level on untreated controls = 85%

Therefore banded mean disease level on untreated controls = 90

$$\begin{aligned} \text{PRCO} &= 100 - \left\{ \frac{\text{Banded mean disease level for treatment A}}{\text{Banded mean disease level on untreated controls}} \right\} \times 100 \\ &= 100 - \left(\frac{30}{90} \times 100 \right) = 66.7 \end{aligned}$$

The PRCO is then rounded to the nearest whole number; therefore, in this particular example, the PRCO result is 67.

It is possible for negative PRCO values to be obtained.

PRCO results are shown below.

TABLE I

COMPOUND NO.	ERYSGT	PUCCRT	PHYTIN
	6hour Protectant	1day Protectant	1day Protectant
(I)	100	100	78


Key to Table I

ERYSGT = *Blumeria graminis tritici* PUCCRT = *Puccinia triticina*


PHYTIN = *Phytophthora infestans lycopersici*

CLAIMS.

1. A compound having the formula (I)



2. A compound according to claim 1 having the formula (II)



3. A compound as claimed in claim 1 or claim 2 in the form of a whole fermentation broth containing such a compound; the solids of a whole fermentation broth containing such a compound; intact or lysed mycelia separated from a whole fermentation broth containing such a compound; the solids of such a whole fermentation broth containing such a compound after separation of intact or lysed mycelia; or such a whole fermentation broth containing such a compound after the separation of solids and the mycelia.
4. A biocidal composition comprising a biocidally effective amount of a compound as claimed in any of the preceding claims.
5. A composition according to claim 4 wherein said composition is a fungicidal composition comprising a fungicidally effective amount of a compound of formula (I) and a suitable carrier or diluent therefor.

6. A method of combating and controlling fungi which comprises treating the fungi or the locus of the fungi with a composition according to claim 5.
- 5 7. A method of combating or controlling phytopathogenic fungi according to claim 6 which comprises applying a fungicidally effective amount of a compound of formula (I), or a composition containing a compound of formula (I), to a plant, to a seed of a plant, to the locus of the plant or seed or to soil or any other plant growth medium, e.g. nutrient solution.
- 10 8. A process for the production of a compound of formula (I) which comprises the step of cultivating an organism of the strain *Streptomyces malaysiensis* capable of producing a compound of formula (I) whereby said compound is produced, and if desired isolating said compound therefrom.
- 15 9. A process according to claim 8 wherein said strain is *Streptomyces malaysiensis* JHCC 553434 (DSM 14702) or a mutant thereof.
- 20 10. A process according to claim 8 or claim 9 in which a compound of formula (I) is separated from the fermentation broth.
- 25 11. *Streptomyces malaysiensis* JHCC 553434 (DSM 14702) and mutants thereof.